

# Seminar\_YISHPESS\_CHERY revisi

*by* Cerika Rismayanthi

---

**Submission date:** 21-Aug-2019 05:00PM (UTC+0700)

**Submission ID:** 1161981989

**File name:** Seminar\_YISHPESS\_CHERY\_revisi.docx (184.28K)

**Word count:** 3347

**Character count:** 17982

# THE PREDICTION OF THE INCIDENCE RATE OF CARDIOVASCULAR DISEASE FOR THE EMPLOYEES AND LECTURERS OF YOGYAKARTA STATE UNIVERSITY BASED ON THE POST-EXERCISE RECOVERY HEART RATE

**Prijo Sudibjo dan Cerika Rismayanthi**  
**Health and Recreation Education Departement**  
**Yogyakarta State University**

**Background:** Work productivity is highly demanded in ensuring the sustainability of an institution. The fitness levels and health levels greatly affect a person's work productivity. There are several fitness indicators, one of which is cardiovascular fitness. The Rockport test is a cardiovascular fitness test that can be performed and has a high level of safety. To predict the level of cardiovascular health can be done with heart rate recovery measurements.

**Objective:** The purpose of this study were to determine: (1) the fitness levels of Yogyakarta State University lectures and employees, and (2) the magnitude of risk for the occurrence of cardiovascular disease.

**Methods:** This study is a cross sectional descriptive study on Yogyakarta State University lectures and employees who volunteered to take a fitness test by Department of Health Education and Recreation FIK UNY. The subject of this research amounted to 63 people. All subjects were given a fitness test of the rockport method and measured recovery heart rate after fitness test. From the result of recovery heart rate obtained then determine the magnitude of risk for the occurrence of cardiovascular disease. Data was presented descriptively qualitatively by percentage.

**Results:** The result of this study indicate that: (1) the fitness level of total 63 subjects, showed 31 (49%) subjects has very low fitness level, 10 (16%) subjects has low fitness level, 20 (32%) subjects has medium fitness level, and 2 (3%) subjects has good fitness level, (2) the results of the study also indicate that 37 (58,73%) subjects has high risk of cardiovascular disease, and 26 (41,27%) subjects has low risk of cardiovascular disease.

**Conclusions:** Measurements of fitness level and prediction of cardiovascular disease occurrence on subjects showed that the majority of subjects were 31 (49%) subjects has low fitness level, and 37 (58, 73%) subjects has a high risk of cardiovascular disease.

*Key Words: The Incidence Rate Of Cardiovascular Disease, Post-Exercise Recovery Heart Rate*

## **Background**

Every individual who works for an institution, either government and private institution, is always required to have maximum work productivity. Maximum work productivity will be achieved if a person has an optimal level of health and fitness. On the other hand, due to the advancement of technology that will facilitate every activity of a person, it will encourage a person to be inactive by trying to reduce his/her physical activities which will eventually lead to decreased

fitness and even trigger the incidence of diseases. One of the diseases that is closely related to the level of physical activity of a person is cardiovascular disease, such as, heart disease, hypertension, and stroke.

In general, fitness is defined as the ability to perform daily activities efficiently without causing excessive fatigue, so that a person can still enjoy his/her spare time. Giriwijoyo and Sidik (2012: 20-21) said that fitness is a state of physical ability that can adjust the function of the body to certain physical tasks and/or to the environmental condition that must be overcome efficiently, without excessive fatigue, and has been perfectly restored before performing the same tasks in the following day. There are several components of fitness, one of which is cardiovascular fitness or heart and lung fitness that describes the ability of the lungs and heart to meet the oxygen demand for muscles' work for a long time. It can only be achieved by performing regular and programmed exercise.

The more fit a person, the more efficient the work of heart in fulfilling the blood circulation to all tissues and organs of the body. It can be seen by the decreasing frequency of resting heart rate which is close to 60 beats per minute. In contrast, the more unfit a person, the faster the heart rate which will approach to 90 beats per minute. Thus, the heart rate is an indicator of a person's fitness that can be easily observed. When performing any physical activity or exercise, the body will strive to meet the needs of oxygen and energy for muscle contraction by increasing heart rate. This reaction is an acute reaction of the body in an effort to meet the needs for muscle contraction. However, the long-term effects of the regular and programmed exercise will improve the efficiency of cardiovascular work as the stroke volume increased as a result of the left ventricular hypertrophy.

As with the measurement of body composition and muscle strength, the measurement of cardiac capacity is also important in assessing the performance of a person or overall cardiovascular fitness. There are several heart rate measurements that can be used to monitor a person's health, such as, resting heart rate, maximum heart rate, *reserve heart rate*, and *recovery heart rate*. *Recovery heart rate* (RHR) is a measurement of heart rate after performing aerobic physical exercise. It is well known that RHR is one of the best indicators that can assess both overall and cardiovascular health conditions (Stone, 2011: 1-2).

The fitness of a person can be identified by conducting an objective and standardized fitness test, which is by measuring the status of fitness components that includes the measurements of heart and lung resistance, muscle strength and endurance, flexibility, and body fat percentage. Physical fitness test is a procedure in the form of physical exercises which is aimed at measuring fitness level according to the purpose. The level of physical fitness can be known by physical fitness test and measurement. In the measurement test, there are several principles and procedures which have to be performed. The Rockport method is one of the safest methods of heart and lung fitness measurements for all ages, especially for older people. This method is also safe for groups of people suffer from certain diseases. In addition, the implementation of the Rockport method also does not require for medical examination before performing the fitness test (Adams, 2002: 139).

The heart rate is intrinsically regulated from the heart, both autonomously and extrinsically, as a response to physical and mental activities and is also influenced by the autonomous and hormonal nervous system. The reflect of heart rate regulation is related to the excitatory input from the cardiorespiratory and baroreceptor systems in the blood vessels. The <sup>1</sup> heart rate describes the dynamic balance between the sympathetic and parasympathetic nervous system. The

parasympathetic activation inhibits the heart rate by acetylcholine release from efferent fibers of vagus nerve. While the sympathetic activation will trigger the heart rate by releasing either epinephrine in the circulation or norepinephrine in the nervous system, or both. (Robertson, *et al.*, 1979).

The abnormality of autonomic nervous system is closely related to the increased mortality and morbidity of cardiovascular disease in both healthy populations and populations suffering from certain diseases. There are many markers of cardiovascular activity used clinically in identifying patients at high risk for sudden cardiac death. One of the markers is the *exercise stress test* which is used as the diagnostic and prognostic tool that is used extensively in evaluating cardiovascular function. It is known that the exercise response is acutely influenced by the activity of autonomic nervous system, where there is increased sympathetic and decreased parasympathetic activities while performing a physical activity. While after performing an activity, there is a combination of decreased sympathetic activity and parasympathetic reactivation. Thus, Recovery HR after performing the maximum physical activity is one of the simplest parameters that can describe the autonomic activity of the heart and can predict the presence of cardiovascular disorders. (Cole *et al.*, 1999: 1351-57 and Jouven *et al.*, 2005: 1951-58).

It is known that there are several types of heart rates that can be used for monitoring the health and fitness levels, such as, resting heart rate, maximal heart rate, *heart rate reserve*, and *recovery heart rate*. There are several ways in measuring the heart rate. Resting heart rate is calculated in the morning when awakened from sleep before performing any activity, expecting that the cardiac output at that time is reserved for basal metabolism without being affected by either physical or mental activity (Stone, 2011:2). Normal heart rate ranges from 60 – 100 beats per minute. The heart rate less than 60 beats per minute is referred to bradycardia and which is more

than 100 beats per minute is referred to tachycardia, both of which are the conditions of cardiovascular abnormalities.

The maximum heart rate can be determined by measuring the heart rate immediately after performing a very heavy exercise. However, it is very dangerous to do by someone who does not have a good level of health and should be under the supervision of a doctor. To avoid this risk, there has been many ways that are used recently for measuring the maximum heart rate. There are several calculations to estimate the maximum heart rate by ages, among others, as follows: (Stone, 2011:2)

- 220 - Age (th) for Men, and
- 226 - Age (th) for Women

The *heart rate reserve/HRR* is the difference between the maximum heart rate and resting heart rate, or it can be calculated by using the following formula:

$$\text{HRR} = \text{HR max} - \text{HR rest}$$

The greater the number of HRR, the higher the fitness of a person, which also means that the lower the HR rest, the more excellent the fitness of a person. (Stone, 2011:2)

*Recovery Heart Rate (RHR)* is a measurement of heart rate after performing aerobic exercise. This measurement is performed during the cooling-down period, so that it can monitor changes in heart rate back to normal after performing an exercise. RHR measurement can be performed by measuring the heart rate immediately after performing an exercise and measuring the heart rate two minutes after the exercise. RHR is the difference between the two heart rate calculations (Stone, 2011:2).

It is known that increased heart rate during physical exercise and decreased heart rate after the exercise is a role of the change to the sympathetic and parasympathetic nervous systems. The slow decrease in RHR after performing any physical exercise is associated with higher mortality, cardiac death, and incidence rate of cardiovascular disease associated with ischemic processes (Ghaffari et.al.,2011:47-54). Based on the formulation of the problems above, the objectives of this study are to determine the fitness level of the lecturers and employees of Yogyakarta State University, and to find out the level of risk against the incidence of cardiovascular disease.

### **RESEARCH METHOD**

This study is a *cross-sectional* descriptive study conducted on lecturers and employees of Yogyakarta State University who voluntarily took a fitness test held by the Department of Health Education and Recreation of Yogyakarta State University. The subjects of study were taken by conducting *purposive sampling*. All subjects were given for a Rockport fitness test and measured for their *recovery heart rate* (RHR) after performing a fitness test. The RHR measurement was obtained by subtracting the maximum heart rate after performing a Rockport test with a heart rate two minutes after the completion of Rockport test. The RHR is abnormal if the  $RHR \leq 12$  beats. From the *recovery heart rate* obtained, the level of risk was then able to determine regarding the incidence of cardiovascular disease. There were also anthropometric measurements conducted in the form of body height, abdominal circumference, and hip circumference measurements. The whole data would be presented descriptive-qualitatively in a percentage.

### **RESULTS AND DISCUSSION**

The characteristics of the subjects of study were lecturers and employees of Yogyakarta State University who voluntarily took a fitness test held by the Department of Health Education and Recreation, which were briefly presented as follows: the subjects consisted of 50 men (79%) and

13 women (21%). The distribution of the subjects of this study by sex was dominated by men. The majority of subjects in this study were in a late-adolescence age of 17-25 years old as many as 46 people (73%), while the characteristics of the subjects based on physical fitness level with a very poor level were 31 people (49%). The data characteristics of the subjects can visually be seen in the following diagram:

Data characteristic of the research subject can be visually seen in the following diagram:



Figure 1. Characteristics of Research Subjects by Sex





Figure 2. Characteristics of Research Subjects Based on Age Distribution



Figure 3. Characteristics of Research Subjects by Level of Fitness

## Results

The assessment of the prediction of incidence rate of cardiovascular disease in this study was based on the results of measurement of *recovery heart rate (RHR)* of the subjects. All subjects were given for a Rockport-method fitness test and measured for their *recovery heart rate (RHR)* after performing the fitness test. The RHR measurement was obtained by subtracting the maximum heart rate after performing a Rockport test with a heart rate two minutes after the completion of Rockport test. The *RHR* is abnormal if the  $RHR \leq 12$  beats. The results of data analysis can be seen in table 4.2 below:

**Table 4.2 The Results of Analysis of the Prediction of Incidence Rate of Cardiovascular Disease**

No.	Cardiovascular Disease Category	Norm/Interval Category	Frequency	
			f (n)	%
1.	High Risk	$\leq 12$	37	58,73
2.	Low Risk (Normal)	12	26	41,27
<b>Total</b>			<b>63</b>	<b>100</b>

Based on the distribution of frequency above, it can be seen that 37 (58,73%) of 63 employees of Yogyakarta State University, as the subjects of this study, have a high risk of cardiovascular disease, and 26 employees (41,27%) have a low risk of cardiovascular disease. The data shows that the majority of the subjects has a high risk of having cardiovascular disease.

From the table results of prediction analysis of the incidence rate of cardiovascular disease, can be visually presented in the form of histogram as follows:




Figure 4. Histogram Prediction of Cardiovascular Disease Occurrence Rate at Employees State University of Yogyakarta

### Discussion

Nowadays, the mortality and morbidity rates of cardiovascular disease are increasing. The abnormality of autonomic nervous system is closely related to the increasing mortality and morbidity rates of cardiovascular disease in both healthy populations and populations suffering from certain diseases. There are many markers of cardiovascular activity used clinically in identifying patients at high risk for sudden cardiac death. It is known that the exercise response is

acutely influenced by the activity of autonomic nervous system, where there is increased sympathetic and decreased parasympathetic activities while performing a physical activity. While after performing an activity, there is a <sup>1</sup> combination of decreased sympathetic activity and parasympathetic reactivation. Thus, Recovery HR after performing the maximum physical activity is one of the simplest parameters that can describe the autonomic activity of the heart and can predict the presence of cardiovascular disorders. (Cole et al, 1999: 1351-57 and Jouven et al, 2005: 1951-58).

The prediction of incidence rate of cardiovascular disease in this study was accurate as based on the *recovery heart rate* measurement. A previous study showed that the patients with abnormal *RHR* showed the decreased parasympathetic activity after performing maximum exercise and abnormal autonomic function (Gayda et.al, 2012: 6). In addition, Stone (2011:1-2) stated that *RHR* is one of the best indicators that can assess both overall and cardiovascular health conditions. The results of data analysis showed that 37 people (58,73%) of the subjects have  $RHR < 12$  which means that the majority of the subjects, which were the employees of Yogyakarta State University, have a high risk of having cardiovascular disease.

The condition of the majority of the subjects who have a high risk of cardiovascular disease can be caused by many factors. One of the factors is the physical activity which has an effect on the fitness level of a person. The fitness level of the subjects showed that the majority of the subjects are in the very poor level which was nearly a half (49%) of the subjects, so that it has an effect on the *RHR* of the subjects. Stone (2012: 2) stated that the greater the number of *HRR*, the higher the fitness of a person, which also means that the lower the *HR rest*, the more excellent the fitness of a person. It can be concluded that the routine physical activity can have an effect on the level of risk of having cardiovascular disease. It is known that increased heart rate during physical

exercise and decreased heart rate after the exercise is a role of the change to the sympathetic and parasympathetic nervous systems. The slow decrease in RHR after performing any physical exercise is associated with higher mortality, cardiac death, and incidence rate of cardiovascular disease associated with ischemic processes (Ghaffari et.al.,2011:47-54).

Dimopoulos *et.al.*(2015:1-10) stated that the activity or position after performing a physical activity has an effect on Recovery HR. Recovery HR  $\leq$  12 beats per minute in the first minute in standing/upright position or  $\leq$  18 beats per minute in the first minute in laying/supine position or  $\leq$  22 beats per minute in the first minute in sitting position shows the abnormal condition. There are several evidences which show that the recovery phase after performing a physical activity can be used as the prognostic factor of the presence of cardiovascular disorders.

### **Conclusions**

Based on the results and discussion presented in the previous chapter, there are some conclusions as follows: The results also showed that 37 subjects (58,73%) have a high risk of having cardiovascular disease, and 26 subjects (41,27%) have a low risk of having cardiovascular disease. In measuring the fitness level (VO2 max) the test must be created in such a way that the pressure on the supply of oxygen to the heart muscle should last a maximum. This Lung Heart Fitness Activity by using ROCKPORT method with run / walk as far as 1.6 km. Study of normality analysis Fitness Level Level of fitness (test rockport) for employees and lecturers Universitas Negeri Yogyakarta Prodi Ikor get fitness test norms that can be in accordance with the conditions of the people of Indonesia so it is expected that a lecturer or employee UNY able to work with productive, efficient and not easily attacked by disease , eager to perform optimally, and tough in completing tasks. Optimum physical fitness can be obtained through proper physical practice, regular, and measurable. With fitness test,

employees at Yogyakarta State University get the chance to know the norm and profile of health and fitness level, so that by knowing the fitness level of lecturer and employee of UNY can do special intervention if found that fitness level is not optimal.

### **Suggestions**

Based on the conclusions above, it is suggested for the employees of Yogyakarta State University to improve their physical activities or exercises and to perform exercises measurably and regularly.

### **References**

- Adams, G.M., 2002, Exercise Physiology, Laboratory Manual, Fourth Edition, The McGraw-Hill Companies Inc., New York.pp: 139-141.
- Cole CR, Blackstone EH, Pashkow FJ, Snader CE, Lauer MS., 1999. Heart-Rate Recovery Immediately After Exercise as a Predictor of Mortality. N Eng J Med. Vol. 341, pp: 1351-1357.
- Dimopoulos S, Manetos C, Panagopoulou N, Karatzanos Land Nanas S., 2015.The Prognostic Role of Heart Rate Recovery after Exercise in Health and Disease, Austin J Cardiovasc Dis Atherosclerosis, Volume 2 Issue 2 -2015, pp:1-10
- Gayda, M. et al. 2012. Heart Rate Recovery After Exercise and Long-term Prognosis in Patients With Coronary Artery Disease, Canadian Journal of Cardiology. Vol, 20, pp. 1-7.
- Ghaffari, S., Kazemi, B, Aliakbarzadeh, P.2011. Abnormal Heart Rate Recovery after Exercise Predicts Coronary Artery Disease Severity, Cardiology Journal, Vol. 18, No. 1, pp. 47-54.
- Giriwijoyo, S., dan Sidik, D.Z., 2012, Ilmu Faal Olahraga (Fisiologi Olahraga), Fungsi Tubuh Manusia pada Olahraga Untuk Kesehatan, PT Remaja Rosdakarya Offset, Bandung.pp: 20-21.
- Jouven X, Empana JP, Schwartz PJ, Desnos M, Courbon D, Ducimetière P., 2005. Heart-rate profile During Exercise as a Predictor of Sudden Death. N Eng J Med.; Vol. 352, pp: 1951-1958.
- Robertson D, Johnson GA, Robertson RM, Nies AS, Shand DG, Oates JA., 1979. Comparative assessment of stimuli that release neuronal and adrenomedullary catecholamines in man. Circulation.; Vol. 59, pp: 637-643.

Stone, J. 2011. Cardiovascular: HR rest, HR max, HRR, HR Recovery and VO2 Max., John Stone Fitness, <http://www.johnstonefitness.com/2011/02/14/cardiovascular-fitness-hrrest-hrmax-hrr-recovery-hr-and-vo2-max/>.pp:1-2

# Seminar\_YISHPESS\_CHERY revisi

---

## ORIGINALITY REPORT

---

**2%**

SIMILARITY INDEX

**2%**

INTERNET SOURCES

**0%**

PUBLICATIONS

**0%**

STUDENT PAPERS

---

## PRIMARY SOURCES

---

**1**

**austinpublishinggroup.com**

Internet Source

**2%**

---

Exclude quotes      On

Exclude matches      < 2%

Exclude bibliography      On

# Seminar\_YISHPESS\_CHERY revisi

---

## GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---

PAGE 11

---

PAGE 12

---

PAGE 13

---